

兔抗 MOK 多克隆抗体

中文名称: 兔抗 MOK 多克隆抗体

英文名称: Anti-MOK rabbit polyclonal antibody

别名: RAGE; RAGE1; STK30; RAGE-1

储 存: 冷冻(-20℃) 避光

抗 原: MOK

宿 主: Rabbit

反应种属: Human

相关类别: 一抗

标记物: Unconjugate

克隆类型: Unconjugate

技术规格

Background:	This gene belongs to the MAP kinase superfamily. The gene was found to be regulated by caudal type transcription f actor 2 (Cdx2) protein. The encoded protein, which is localized to epithelial cells in the intestinal crypt, may play a role in growth arrest and differentiation of cells of upper crypt and lower villus regions. Multiple alternatively spliced transcript variants encoding different isoforms have been observed for this gene.
Applications:	WB, IHC
Name of antibody:	MOK
Immunogen:	Synthesized peptide derived from internal of human MOK.
Full name:	MOK protein kinase



www.shjning.com

Synonyms:	RAGE; RAGE1; STK30; RAGE-1
SwissProt:	Q9UQ07
IHC positive control:	Human brain tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	48 kDa
WB Positive control:	HepG2 cells, COLO205 cells and 293 cells lysates
WB Recommended dilution:	500-3000





