

兔抗 TCL1A 多克隆抗体

中文名称: 兔抗 TCL1A 多克隆抗体

英文名称: Anti-TCL1A rabbit polyclonal antibody

别 名: TCL1A; TCL1

相关类别: 一抗

存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: TCL1A

反应种属: Human, Mouse, Rat

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	TCL1 (T cell leukemia 1), MTCP1 and TCL1b belong to the TCL1 proto-oncogene family, and their products are involved in Akt activation during embryonic development, T cell leukemias, prolymphocytic leukemias and B cell lymphomas. The Akt association domain of TCL1 binds with the PH domain of Akt. The formation of an oligomeric TCL-Akt complex is required for TCL1 coactivator function and results in phosphorylation and activation of Akt. Furthermore, functional analysis indicates that the interaction between TCL1 and Akt promotes translocation of Akt to the nucleus. The se findings are supported by the crystal structure of TCL1, which suggests that TCL1 may participate in molecular transport.
Applications:	WB
Name of antibody:	TCL1A



www.shjning.com

Immunogen:	Fusion protein of human TCL1A
Full name:	T-cell leukemia/lymphoma 1A
Synonyms:	TCL1A; TCL1
SwissProt:	P56279
WB Predicted band size:	13 kDa
WB Positive control:	Ramos cells
WB Recommended dilution:	500-2000

