

## BCL10 抗原（重组蛋白）

中文名称： BCL10 抗原（重组蛋白）

英文名称： BCL10 Antigen (Recombinant Protein)

别名： CLAP; mE10; CIPER; c-E10; CARMEN

储存： 冷冻（-20℃）

相关类别： 抗原

### 概述

Full length fusion protein

### 技术规格

<b>Full name:</b>	B-cell CLL/lymphoma 10
<b>Synonyms:</b>	CLAP; mE10; CIPER; c-E10; CARMEN
<b>Swissprot:</b>	O95999
<b>Gene Accession:</b>	BC053617
<b>Purity:</b>	>85%, as determined by Coomassie blue stained SDS-PAGE
<b>Expression system:</b>	Escherichia coli
<b>Tags:</b>	His tag C-Terminus, GST tag N-Terminus
<b>Background:</b>	This gene was identified by its translocation in a case of mucosa-associated lymphoid tissue (MALT) lymphoma. The protein encoded by this gene contains a caspase recruitment domain (CARD), and has been shown to induce apoptosis and to activate NF-kappaB. This protein is reported to interact with other CARD domain containing proteins including CARD9, 10, 11 and 14, which are thought to function as upstream regulators in NF-kappaB signaling. This protein is found to form a complex with MALT1, a protein encoded by another gene known to be translocated in MALT lymphoma.

